

Phenylpropanoid Glucosides from *Eucalyptus maculata*

Omar A. Rashwan*

Pharmacognosy Department, Faculty of Pharmacy, Cairo University, Cairo 11562, Egypt.

* To whom correspondence should be addressed; e-mail

Received: 25 October 2024; in revised form 23 December 2024 / Accepted: 24 December 2024 /
Published: 30 January 2024

Abstract: Three compounds were isolated from the butanol soluble fraction of the resinous exudate from the stem of *Eucalyptus maculata*. In addition to *p*-coumaric acid two new compounds were identified. They were identified as 1-*O*-cinnamoyl 6-*O*-*p*-coumaroylglucose and 7-methyl-aromadendrin-4'-*O*-(6''-*trans-p*-coumaroyl)- β -D-glucopyranoside by spectroscopic and chemical means.

Keywords: *Eucalyptus maculata*, Myrtaceae, diacylglycerol, acyl flavanone glucoside, coumaric acid, NMR, MS.


Introduction

Eucalyptus maculata Hook (Myrtaceae) is a medicinal plant traditionally used for the treatment of asthma and chronic bronchitis [1]. In a recent communication, Abdel-Sattar *et al.* reported the isolation of 1,6-*O*-dicinnamoyl glucose, 7-methylaromadendrin and sakuranetin from the resinous exudate of the stem of this plant [2]. The isolation of a phenylpropanoid from the stem exudate of *E. maculata* [2] was encouraging enough to pursue further investigation of the extracts of this plant. Accordingly, we carried out chemical examination of the butanol soluble fraction of the resin exudate, from which three compounds were isolated.

Results and Discussion

Three phenolic compounds (**1-3**) were isolated from the butanol soluble fraction of the resinous material obtained from the stem of *E. maculata*. Compound **1** was identified as *p*-hydroxycinnamic acid (*p*-coumaric acid) by comparison of its spectral data (MS, NMR) with data reported in the literature [3, 4] and by comparison with an authentic sample (TLC, mp and IR).

Compound **2** gave a positive test for sugars and/or glycosides. HR-FABMS gave $[M-H]^-$ at 455.1331, corresponding to the molecular formula $C_{24}H_{24}O_9$ (calculated 455.1342) and this was further confirmed from the ion peak at m/z 479 $[M+Na]^+$ in the electrospray ionization (ESI) mass spectrum (positive mode). TLC examination of the acid and alkaline hydrolysates revealed the presence of cinnamic acid, *p*-coumaric acid, as well as glucose as the sugar moiety. The β -configuration of the glucose moiety was indicated from the large coupling constant of its anomeric proton (cf. Agrawal, [5]), and by the downfield shift (δ 5.53) of this proton [6]. The previous finding was also supported by the similarity of 1H - and ^{13}C -NMR spectral data of compound **2** with those of 1,6-dicinnamoyl- O - α -D-glucopyranoside (**3**), previously isolated from the same plant [2].



The presence of a cinnamoyl moiety in **2** was confirmed by comparison of its NMR data with those of **3**. However, one of the cinnamoyl moieties in **3** was replaced by a *p*-coumaroyl moiety in **2**, as was ascertained from the observation of proton signals due to a *p*-disubstituted aromatic ring system [δ_{H} 7.55 (2H, d, J = 8.5 Hz, H-2, H-6) and 6.80 (2H, d, J = 8.5 Hz, H-3, H-5)]. Acylation at C-1 and C-6 of the glucose unit was confirmed by comparison of the ^1H - and ^{13}C -NMR chemical shifts of the glucose unit compared with those reported in **3** [1], which suggested acylation, as in **3**. The acylation at C-6 of the glucose moiety was determined from the downfield shift of the C-6 (+ 2.4) relative to the respective carbon in β -glucopyranose [5]. TLC of the partial acid hydrolysate of **2** using 5% HCl [7] indicated that *p*-coumaroyl moiety was located at C-1 (*p*-coumaric acid was first identified on TLC) and cinnamoyl moiety was located at C-6 (cinnamic acid was later identified). This finding was further supported from the HMBC spectrum, which showed long-range correlation between the carbonyl of the coumaric acid at δ_{C} 166.8 to the anomeric proton (δ_{H} 5.53) and to the protons at δ_{H} 6.65 and 7.65 (H_{α} and H_{β} , respectively). From the foregoing findings, compound **2** was thus identified as 1-*O*-cinnamoyl 6-*O*-*p*-coumaroyl- β -D-glucopyranoside, which represents a new natural product described here for the first time.

Compound **4** was obtained as white amorphous powder, $[\alpha]_{\text{D}}^{25} + 29.4^{\circ}$ (c. 0.2, MeOH). HR-FABMS of compound **4** gave $[\text{M}-\text{H}]^{-}$ at 609.1658, assigned to the molecular formula $\text{C}_{31}\text{H}_{30}\text{O}_{13}$ (calculated 609.1667). UV spectral analysis of **4** in MeOH (220, 292, 310_{sh} nm) and after addition of different shift reagents suggested the presence of a flavanonol skeleton with free hydroxyl groups at 3 and 5 positions [8]. The ^1H -NMR spectrum of **4** showed the presence of two doublets at δ_{H} 4.18 and 5.86 (J =11.6 Hz), characteristic of the *trans* H-2/H-3 protons, and supported the above finding. By acid hydrolysis, the flavonoid aglycone part was identified as 7-*O*-methyl aromadendrin, previously isolated from the same plant [2]. The results of acid and alkaline hydrolysis revealed in addition to 7-*O*-methyl aromadendrin, the presence of *p*-coumaric acid and glucose (TLC). The coupling constant of the anomeric proton indicated the β -configuration of the glucose moiety [5]. The glycosylation position was determined to be at C-4' hydroxyl group from the observation of the downfield shift of C-4' (δ 157.6) and the long-range correlation between H-1'' and C-4' in the HMBC spectrum of **4**. Acylation of glucose moiety at C-6'' was established from the downfield shift of C-6'' (+ 1.4 ppm) and the upfield shift of C-5'' (- 2.7 ppm) relative to the respective carbons in glucopyranose [5]. From the aforementioned data, compound **4** was elucidated as 7-methylaromadendrin-4'-*O*-(6''-*trans*-*p*-coumaroyl)- β -D-glucopyranoside, which was isolated here for the first time from the title plant and from nature.

Acknowledgments

The author is indebted to Dr. Essam Abdel-Sattar and Dr. Meselhy R. Meselhy, Department of Pharmacognosy, College of Pharmacy, Cairo University, Egypt, for running the NMR and MS for their valuable discussions and to Prof. Dr. Mahmoud A. M. Nawwar for running the 270 MHz NMR spectra.

Acid hydrolysis of 4.

The hydrolysis was performed according to ref. [7, 8].

1-O-Cinnamoyl 6-O-coumaroyl- β -D-glucopyranoside (2).

Molecular formula $C_{24}H_{24}O_9$, $[\alpha]_D^{25}$ - 24.6 ° (c. 0.35, MeOH), IR $\nu_{\max}$ (KBr) cm^{-1} : 3500, 1720, 1660, 1605, 830. UV (MeOH) $\lambda_{\max}$ (ϵ): 286 (2.8×10^3), 312 (2.3×10^3) nm. 1H -NMR (270 MHz, CD_3OD) δ : **cinnamoyl**: 7.75 (2H, m, H-2, H-6), 7.65 (1H, d, $J = 16$ Hz, H_β), 7.40 (3H, m, H-3, H-4, H-5), 6.40 (1H, d, $J = 16$ Hz, H_α), **p-coumaroyl**: 7.65 (1H, d, $J = 16$ Hz, H_β), 7.55 (2H, $J = 8.5$ Hz, H-2, H-6), 6.80 (2H, d, $J = 8.5$ Hz, H-3, H-5), 6.65 (1H, d, $J = 16$ Hz, H_α), **β -D-glucose**: 5.53 (1H, d, $J = 7$ Hz, H-1), 4.45 (1H, d, $J = 11$ Hz, H-6A), 4.20 (1H, dd, $J = 11, 4.5$ Hz, H-6B), 3.60 (1H, m, H-5), 3.40-3.55 (3H, m, H-2, H-3, H-4). ^{13}C NMR (67.5 MHz, CD_3OD) δ : **cinnamoyl**: 167.0 (C=O), 145.6 (C_β), 135.2 (C-1), 129.7 (C-2, C-4, C-6), 129.1 (C-3, C-5), 118.6 (C_α), **p-coumaroyl**: 166.0 (C=O), 160.8 (C-4), 146.8 (C_β), 131.1 (C-2, C-6), 126.7 (C-1), 116.6 (C-3, C-5), 114.7 (C_α), **β -D-glucose**: 95.1 (C-1), 77.6 (C-3), 75.7 (C-5), 73.8 (C-2), 70.8 (C-4), 64.2 (C-6). HR-MS $[M-H]^-$ (negative mode) m/z 455.1331 (calculated 455.1342). ESI-MS (positive mode) m/z (rel. int.): 479 $[M + Na]^+$ (100), 293 $[M - p\text{-coumaroyl}]^+$ (31), 131 $[Ph-C=C-CO]^+$ (55).

7-Methylaromadendrin-4'-O-(6''-trans-p-coumaroyl)- β -D-glucopyranoside (4)

Obtained as a white amorphous powder. $[\alpha]_D^{25}$ + 29.4 ° (c. 0.2, MeOH), IR $\nu_{\max}$ (KBr) cm^{-1} : 3448, 2922, 1686, 1637, 1510, 1075 and 833 cm^{-1} ; UV (MeOH) $\lambda_{\max}$ ($\log \epsilon$): 220 (4.35), 293 (4.26), 315 (4.20); (+ NaOMe): 220, 245_{sh}, 292, 362, $AlCl_3$ 222, 314, 370; (+ $AlCl_3/HCl$): 222, 314, 370; (+ NaOAc): 222, 293, 315_{sh}; (+ NaOAc/ H_3BO_3): 224, 292, 315_{sh}. 1H -NMR (400 MHz, CD_3OD): **flavanonol**: δ 3.74 (3H, s, OMe), 4.18 (1H, d, $J = 11.6$ Hz, H-3), 5.86 (1H, d, $J = 11.6$ Hz, H-2), 5.93 (1H, br d, H-6), 6.02 (1H, br d, H-8), 7.03 (2H, d, $J = 8.8$ Hz, H-3', H-5'), 7.30 (2H, d, $J = 8.8$ Hz, H-2', H-6'), **p-coumaroyl**: δ 7.56 (1H, d, $J = 16$ Hz, H_β), 7.32 (2H, d, $J = 8$ Hz, H-2'', H-6''), 6.67 (2H, d, $J = 8$ Hz, H-3'', H-5''), 6.62 (1H, d, $J = 16$ Hz, H_α), **β -D-glucose**: δ 4.83 (1H, d, $J = 6.5$, H-1''), 4.46 (1H, d, $J = 11$, H-6_A''), 4.20 (1H, dd, $J = 11, 4.5$, H-6_B''), 3.60 (1H, m, H-5''), 3.40-3.55 (3H, m, H-2'', H-3'', H-4''). ^{13}C NMR (100 MHz, CD_3OD) **flavanonol**: δ 196.3 (C-4), 168.3 (C-7), 163.3 (C-9), 162.5 (C-5), 157.6 (C-4'), 130.4 (C-1'), 128.6 (C-2', C-6'), 116.6 (C-3', C-5'), 101.0 (C-10), 95.8 (C-8), 94.1 (C-6), 82.9 (C-2), 72.2 (C-3), **β -D-glucose**: δ 100.6 (C-1''), 76.3 (C-3''), 74.0 (C-5''), 73.0 (C-2''), 70.1 (C-4''), 63.2 (C-6''), **p-coumaroyl**: δ 167.6 (C=O), 159.4 (C-4'''), 145.5 (C_β), 129.8 (C-2''', C-6'''), 125.5 (C-1'''), 115.6 (C-3''', C-5'''), 113.7 (C_α). HR-MS $[M-H]^-$ (negative mode) m/z 609.1658 (calculated 609.1667). ESI-MS m/z (rel. int.): 461 $[M - p\text{-coumaroyl}]^+$ (28), 325 $[M - \text{methylaromadendrinoyl} + Na]^+$ (10), 259 (75), 223 (44), 149 (100).

References

1. Mishra, C. S. and Misra, K. *Planta Medica* **1980**, 38, 169.
2. Abdel-Sattar, E., Kohiel, M. A., Shehata, I. A. and El-Askary, H. *Pharmazie* **2000**, 55, 263.
3. Chaudhuri, P. K. and Thakur, R. S. *Phytochemistry* **1986**, 25, 1770.
4. Itokawa, H., Suto, K. and Takeya, K. *Chem. Pharm. Bull.* **1981**, 29, 254.
5. Agrawal, P. K. *Phytochemistry* **1992**, 31, 3307.
6. Latza, S., Gansser, D. and Berger, R. G. *J. Agr. Food Chem.* **1996**, 44, 1367.
7. Nawwar, M. A. M. and Hussein, S. A. M. *Phytochemistry* **1994**, 36, 1035.
8. Mabry, T. J., Markham, K. R. and Thomas, M. B.: *The Systematic Identification of Flavonoids*. Springer: Berlin, 1970; 35.

Sample Availability: Available from the authors.

© 2002 by MDPI (<http://www.mdpi.org>). Reproduction is permitted for noncommercial purposes.